

Patient information from BMJ

Last published: Jul 29, 2020

Irritable bowel syndrome: what treatments work?

Irritable bowel syndrome (IBS) is a condition that can cause stomach pain, cramps, bloating, constipation, and diarrhoea. There isn't a cure for IBS, but lifestyle changes and treatments can improve your symptoms.

You can use our information to talk to your doctor and decide which treatments are right for you. To learn more about IBS and its symptoms, see our leaflet *Irritable bowel syndrome: what is it?*

Lifestyle and diet

Keeping a symptom diary

Certain things in your life, such as stress and certain foods, might bring on your IBS symptoms. Your doctor might recommend that you keep a symptom diary.

This will allow you to track when you get symptoms, so you can see whether anything seems to trigger them. Once you identify a possible trigger, you can take steps to avoid it.

Managing stress

Many people get flare-ups of IBS symptoms when they are feeling stressed. So taking steps to manage stress is often one of the first treatments doctors recommend. Things you might try include:

- deep breathing exercises
- meditation
- yoga
- relaxation classes, and
- taking regular exercise.

Talking to a counsellor or joining an IBS support group might also help. Your doctor can help you explore your options.

Irritable bowel syndrome: what treatments work?

Changing your diet

Sometimes people with IBS have difficulty digesting certain foods, which can make their symptoms worse. These foods can include:

- dairy products
- foods that contain a type of sugar called fructose, and
- coffee and other drinks that contain caffeine.

You might need to cut these out of your diet completely.

With the help of a symptoms diary you might be able to identify foods that make your IBS worse. But it's a good idea to check with a doctor or a dietitian before you make any changes to your diet. Avoiding certain foods can make your life difficult, and it's easy to miss out on important nutrients.

Some people find that their IBS improves when they cut out a type of carbohydrate that is difficult for the body to absorb. These carbohydrates are called FODMAPs. You can ask your doctor or dietitian for details on a **low-FODMAP diet**, or look online.

If constipation is one of your IBS symptoms, you might benefit from eating more **fibre**. Fibre is a bulky substance that your body can't digest. The extra bulk helps make your stools large and soft, and helps your bowels work properly. Good sources of fibre in foods include:

- wholegrain or wholemeal foods
- cereals with bran in them
- vegetables, and
- fresh or dried fruits.

You can also get fibre supplements from pharmacies or health food shops. Some examples are ispaghula husk and methylcellulose. Make sure you drink plenty of water if you're taking fibre supplements. Fibre sometimes gives people wind or makes them feel bloated.

Probiotic supplements can help improve IBS in some people. You can buy these as yoghurts or capsules that contain 'friendly bacteria'.

The idea is that these add to the helpful bacteria that live in your bowels and aid in digestion. Research suggests that one type of probiotic, called *Bifidobacterium infantis*, might be particularly helpful in IBS.

Medicines

If lifestyle and diet changes don't do enough to improve your symptoms, medicines can also help. These include:

- anti-spasmodic drugs, which help to control the muscles around your bowel
- medicines for diarrhoea

Irritable bowel syndrome: what treatments work?

- medicines for constipation, and
- antidepressants.

Your medicines might change depending on what symptoms you are having (for example, constipation or diarrhoea).

Anti-spasmodic drugs

Anti-spasmodic drugs can help with IBS, especially if your main symptom is pain. These drugs work by relaxing the wall of your bowel.

People often take anti-spasmodic drugs for a week or so at a time when their IBS gets bad. It's better to take these medicines when your symptoms flare up, rather than to take them every day. Anti-spasmodics can sometimes cause side effects, such as nausea and an irregular heartbeat.

Peppermint oil works in the same way as anti-spasmodic drugs. It might help reduce pain and bloating. You can buy peppermint oil as drops or tablets from a pharmacy. Peppermint oil sometimes causes heartburn as a side effect.

Medicines for diarrhoea

Doctors often recommend a drug called loperamide to treat bouts of diarrhoea in IBS. You can buy it yourself from a pharmacy. Some people get mild side effects from loperamide. These include dizziness, drowsiness, and bloating.

If loperamide doesn't work for you, there are other options. Talk to your doctor about what might be best for you.

Medicines for constipation

If one of your main symptoms is constipation, you might first want to try eating more fibre or taking fibre supplements. If these don't help, you could also try a laxative to help you go to the toilet.

If you aren't able to take laxatives, or if over-the-counter laxatives don't help enough, your doctor might recommend treatments available only on prescription.

Antidepressants

Some antidepressants seem to help with IBS. Your doctor might suggest an antidepressant if you have bad IBS and other treatments haven't helped you.

You need to take an antidepressant every day for it to help. You probably won't feel the benefits for at least two or three weeks.

Antidepressants can cause side effects, including nausea, weight gain, fatigue, and a dry mouth. However, doctors often prescribe antidepressants at a lower dose for IBS than for depression. So there might be less chance of side effects.

Other treatments

Some people with IBS try **hypnotherapy** to help them relax and cope with their symptoms. If you have this treatment, a therapist helps you focus on your symptoms and what you can do about them while you're in a relaxed state.

There's some research to show that hypnotherapy might help with IBS, but it can take up a lot of time and can be expensive.

Cognitive behaviour therapy is a talking treatment (a psychotherapy). It aims to help change the way you think about your illness and give you more control over it. Studies have suggested that it might help people get fewer symptoms of IBS.

The patient information from *BMJ Best Practice* from which this leaflet is derived is regularly updated. The most recent version of Best Practice can be found at bestpractice.bmj.com. This information is intended for use by health professionals. It is not a substitute for medical advice. It is strongly recommended that you independently verify any interpretation of this material and, if you have a medical problem, see your doctor.

Please see BMJ's full terms of use at: bmj.com/company/legal-information. BMJ does not make any representations, conditions, warranties or guarantees, whether express or implied, that this material is accurate, complete, up-to-date or fit for any particular purposes.

© BMJ Publishing Group Ltd 2020. All rights reserved.

